

Phase Diagram of the Two-Channel Kondo Lattice

Mark Jarrell^(a), Hanbin Pang^(a), and D.L. Cox^(b)

^(a) *Department of Physics, University of Cincinnati, Cincinnati, OH 45221*

^(b) *Department of Physics, The Ohio State University, Columbus, OH, 43202*

(February 1, 2008)

The phase diagram of the two-channel Kondo lattice model is examined with a Quantum Monte Carlo simulation in the limit of infinite dimensions. Commensurate (and incommensurate) antiferromagnetic and superconducting states are found. The antiferromagnetic transition is very weak and continuous; whereas the superconducting transition is discontinuous to an odd-frequency channel-singlet and spin-singlet pairing state.

75.30.Mb, 71.27.+a, 75.10.Dg

Introduction. A number of Heavy Fermion materials display highly unusual superconductivity (HFSCs) [1]. It seems unlikely that conventional superconductivity coexists with the strong local Coulomb correlations necessary to enhance the electronic mass. Indeed, the specific heat jump at the transition scales with the normal state specific heat making it clear pairing is between the heavy electrons (with order 100-1000-fold mass enhancement). The strong coulomb correlations present no problem for unconventional superconducting order parameters with either spatial [2] or temporal nodes (so called “odd-frequency pairing”) [3–7]. Such an interpretation is supported by two sets of data: (i) power laws observed in physical properties below the superconducting transition [1,2], which contrast with the activated behavior of the conventional (nodeless) s -wave order of, e.g., aluminum; (ii) the complex superconducting phase diagrams of UPt₃ and U_{1-x}Th_xBe₁₃ (and possibly UBe₁₃ itself, for which the penetration depth displays evidence of a secondary transition) [1,2,8]. The superconductivity can coexist with antiferromagnetism, which is usually commensurate, as in UPt₃, URu₂Si₂, UPd₂Al₃, and UNi₂Al₃ [1], or compete with it as in CeCu₂Si₂ [9]. Finally, at least in UBe₁₃ [10] and CeCu₂Si₂ [9], the superconductivity arises in a normal state which is clearly *not* described as a Fermi liquid. In each of these materials above T_c the linear specific heat rises with decreasing temperature, the resistivity is approximately linear in T , and the residual resistivity at T_c is high (typically 80-100 $\mu - \Omega$ -cm in the best samples of UBe₁₃).

In this paper, we provide the first calculations of the phase diagram for the two-channel Kondo lattice in infinite spatial dimensions. We find second order antiferromagnetic and first order odd frequency superconducting phase transitions. Coexistence of commensurate (and incommensurate) antiferromagnetism with superconductivity is in principle possible. We present several possible routes to account for the multiple superconducting phases observed in real materials. Taken together with the non-Fermi liquid (NFL) paramagnetic phase discussed in previous publications [12], and earlier suggestions that the two-channel lattice may describe UBe₁₃

and other heavy fermion systems [8], our work establishes the two-channel Kondo lattice model as possessing the key ingredients needed to explain heavy fermion superconductivity.

Motivation A possible model for some heavy fermion compounds is the two-channel Kondo lattice which consists of two identical species of non-interacting electrons antiferromagnetically coupled to an adjacent lattice of spin 1/2 Kondo moments [8]. This model displays Non-Fermi liquid behavior because of the overcompensation of the Kondo spins by the conduction electrons, first pointed out in the single site two-channel model by Nozières and Blandin [11] where extended screening clouds form which retain a spin 1/2 character. As a result, the paramagnetic state is degenerate and the excitation spectrum non-Fermi liquid like. This overscreening results in an interchannel pairing mechanism [8] in the two-channel impurity problem which tends to favor spin-singlet channel-singlet *odd frequency* superconducting fluctuations [5–7]. For the lattice, we find that this overscreening generates novel antiferromagnetic superexchange between the Kondo spins which, taken together with the RKKY exchange, favors antiferromagnetism in the lattice model close to half filling of the bands. The nature of the resulting superconducting and magnetic transitions, as well as the competition between them, is the subject of this manuscript. .

Model The Hamiltonian for the two-channel Kondo lattice is

$$H = J \sum_{i,\alpha} \mathbf{S}_i \cdot \mathbf{s}_{i,\alpha} - \frac{t^*}{2\sqrt{d}} \sum_{\langle ij \rangle, \alpha, \sigma} \left(c_{i,\alpha,\sigma}^\dagger c_{j,\alpha,\sigma} + \text{h.c.} \right) - \mu \sum_{i,\alpha,\sigma} c_{i,\alpha,\sigma}^\dagger c_{i,\alpha,\sigma}, \quad (1)$$

where $c_{i,\alpha,\sigma}^\dagger$ ($c_{i,\alpha,\sigma}$) creates (destroys) an electron on site i in channel $\alpha = 1, 2$ of spin σ , $\mathbf{S}_i$ is the Kondo spin on site i , and $\mathbf{s}_{i,\alpha}$ are the conduction electron spin operators for site i and channel α . The sites i form an infinite-dimensional hypercubic lattice. Hopping is limited to nearest neighbors with hopping integral $t \equiv t^*/2\sqrt{d}$; the scaled hopping integral t^* determines the energy unit and

is set equal to one ($t^* = 1$). Thus, on each site the Kondo spin mediates an interaction between the two different channels. This problem is non-trivial, and for the region of interest in which $J > 0$ and $T \ll J$, t^* it is describable only with non-perturbative approaches. Clearly some simplifying method which allows for a solution of the lattice problem in a non-trivial limit is necessary.

Formalism and Simulation. Such a method was proposed by Metzner and Vollhardt [13] who observed that the renormalizations due to local two-particle interactions become purely local as the coordination number of the lattice increases. A consequence is that the solution of most standard lattice models may be mapped onto the solution of a local correlated system coupled to an effective bath that is self-consistently determined. We refer the reader to recent reviews for further details and references on the method [14].

In order to solve the remaining impurity problem, we use the Kondo impurity algorithm of Fye and Hirsch [15], modified to simulate the two-channel problem [16]. We simulated the model for a variety of fillings and values of the magnetic exchange J ($0 < N \leq 1$ for $J = 0.75, 0.625, 0.5, 0.4$). Error bars on measured quantities are less than 6% for the results presented here. A sign problem was also encountered in the QMC process. It becomes severe for large J and also for low fillings since lower temperatures are required to access the physically interesting regime. This precludes a systematic study of the model with $J > 0.8$ and filling $N < 0.5$.

Both one and two-particle properties are natural products of the QMC simulation. To calculate the spin susceptibility, it was necessary to measure the local three-by-three matrix of the susceptibility including both the Kondo spin and conduction band spin fluctuations. This may then be inverted to calculate the associated irreducible vertex function and the corresponding lattice susceptibility in the usual way [17].

The situation for the superconductivity is a bit more complicated. Since there are no charge fluctuations involving the Kondo spin degree of freedom, only the two conduction channels contribute to the pair-field susceptibility. We can then look for pairing instabilities in singlet and triplet channels for both spin and channel spin. Motivated by the result for the impurity model [5,6] we have restricted our attention to interchannel and inter-spin particle-particle diagrams as shown in Fig. 1. It is possible to make two independent combinations of these diagrams $\chi_{\pm}(i\omega_n, i\omega_m) = \chi_{11}(i\omega_n, i\omega_m) \pm \chi_{12}(i\omega_n, i\omega_m)$, from which we construct a quartet of spin and channel, singlet and triplet pair-field susceptibilities given by

$$P_{SsCs} = T \sum_{nm} f_{-}(i\omega_n) \chi_{-}(i\omega_n, i\omega_m) f_{-}(i\omega_m) \quad (2)$$

$$P_{StCt} = T \sum_{nm} f_{+}(i\omega_n) \chi_{+}(i\omega_n, i\omega_m) f_{+}(i\omega_m) \quad (3)$$

$$P_{StCs} = T \sum_{nm} \chi_{-}(i\omega_n, i\omega_m) \quad (4)$$

$$P_{SsCt} = T \sum_{nm} \chi_{+}(i\omega_n, i\omega_m). \quad (5)$$

Here $f_{\pm}(i\omega_n)$ are odd functions of Matsubara frequency used to project out the odd-frequency pairing, and, for example, $P_{SsCs}(T)$ is the spin-singlet channel-singlet pair-field susceptibility.

To determine the form of $f_{\pm}(i\omega_n)$, we employ the pairing matrix formalism of Owen and Scalapino [18]. Here we represent each of $\chi_{\pm}$ in a two-particle Dyson equation and extract the irreducible vertex functions $\Gamma_{\pm}$. The pairing matrices are then

$$M_{\pm}(i\omega_n, i\omega_m) = \sqrt{\chi_{\pm}^0(i\omega_n)} \Gamma_{\pm}(i\omega_n, i\omega_m) \sqrt{\chi_{\pm}^0(i\omega_m)}. \quad (6)$$

where $\chi_{\pm}^0(i\omega_n)$ are the particle-particle diagrams in Fig. 1 without vertex corrections. $f_{\pm}(i\omega_n)$ is the eigenvector corresponding to the dominant eigenvalue of $M_{\pm}$ (that with the largest absolute value),

Results. In this model antiferromagnetism is driven by both RKKY interactions and a novel type of superexchange. The latter arises from hopping between adjacent spin 1/2 screening clouds, whose overall spin is determined by the conduction electrons; the Pauli principle forbids hopping unless neighboring spins in the same channel are antiparallel. As a result, for large J , the superexchange goes as $\sim (t^*)^2/J$. For conduction band fillings close to $N = 1$ both the RKKY (evaluated at nearest neighbor sites) and the superexchange favor antiferromagnetism (the RKKY exchange remains antiferromagnetic until $N \lesssim 0.5$), and an antiferromagnetic transition results, as shown in Fig. 2. Due to screening of the local moments by the conduction spin, the transition is very weak, as measured by the full susceptibility. Specifically, χ_{AF} is not significantly enhanced over the bulk susceptibility χ_F until $T \gtrsim T_N$. However, the f-electron contribution to the susceptibility shows a protracted scaling region. Here screening effects non-linear feedback that reduces the susceptibility exponent γ from the mean field value $\gamma = 1$. γ increases with doping ($N < 1$), and the transition becomes incommensurate as $T_N \rightarrow 0$.

To explore superconductivity, it is necessary to find the frequency form factors $f_{\pm}$, which, as discussed above, are the eigenvectors corresponding to the dominant eigenvalues of the pairing matrices $M_{\pm}$. As the temperature is lowered, the dominant eigenvalue first becomes large (divergent) and negative, and then abruptly switches to a large and positive value at the transition. This happens first in M_{-} , and the corresponding eigenvector of M_{-} is plotted in the inset to Fig. 3. It can be fit quite accurately to the form $T/2\omega_n$ as shown by the solid line, which corresponds to the form factor of Ref. [4]. Thus, we

use $f_{-}(i\omega_n) = T/2\omega_n$ to project out the odd-frequency pair-field susceptibilities shown in Figs. 3 and 4. (Other form factors $f(i\omega_n) = \tanh(T_0/\omega_n)$ and $\text{sign}(\omega_n)$ were tried and produce qualitatively similar results.)

The first transition is found in the spin-singlet channel-singlet pairing combination, as shown in Fig. 3. (Note that this pair state is of necessity even in parity, so that the odd-frequency condition is required to satisfy the Pauli principle, in contrast with the spin-parity selection rules in the single channel case [3,4].) To interpret this result, remember that the inverse pair field susceptibility is proportional to the curvature of the free energy $f(\Delta_{SsCs})$ as a function of the pairing order parameter $1/P_{SsCs} \propto d^2 f(\Delta_{SsCs})/d\Delta_{SsCs}^2$. Thus, if $P_{SsCs} < 0$ a thermodynamic instability of the system is present. The associated transition cannot be continuous, since this requires the free energy in order parameter space to become flat (ie. P_{SsCs} diverges) so that the order parameter may change continuously. Thus, we identify this as a discontinuous transition. Furthermore, if $P_{SsCs}(T^*) = 0$, then T^* is a lower bound to the transition temperature since when $P_{SsCs}(T - T^* = 0^-) = 0^-$ the curvature of $f(\Delta_{SsCs})$ is divergent and negative. I.e. the free energy displays a downward cusp which would compel the order parameter and the free energy to change discontinuously at T^* , which involves an infinite energy at the transition. Hence, the actual transition occurs at a temperature $T_c > T^*$.

Several remarks are in order about this unusual superconductivity: (a) Fig. 3 suggests that the transition is degenerate at the zone center and the zone corner, and in fact P_{SsCs} vanishes simultaneously over the whole zone, and hence this is a *locally* driven transition! This degeneracy may be lifted when the order parameter becomes finite, otherwise, it will be lifted in finite dimensionality as the superfluid stiffness Υ_s (analogous to spin stiffness in a magnet) must vanish in infinite d . For finite d with $\Upsilon_s > 0 (< 0)$, a local free energy minimum will be found at $\vec{q} = 0 (\vec{q} \neq 0)$. (b) The vanishing of the local pair susceptibility here at T^* contrasts with the impurity model in which it diverges logarithmically as $T \rightarrow 0$ [5,6]. (c) The excitation spectrum of such a transition may well be highly exotic [4].

As shown in Fig. 4, the ground state of the system may be superconducting or magnetic which may coexist or compete. Our work gives only estimates of the transition temperatures; detailed exploration within the ordered phases will answer this question definitively. In general superconductivity will occur first (at the highest transition temperature) away from half filling and antiferromagnetism will occur first near half filling. However, for values of $J > 0.75$, we find that superconductivity occurs first even at half filling (the minus sign problem precluded a systematic study for these values of J so they are not presented in Fig. 4). We also find that, $T^* \propto T_0$, with the constant of proportionality ≈ 0.5 weakly depen-

dent on J ; whereas T_N appears to depend upon both J and T_0 .

Speculation and Interpretation We offer two compatible interpretations of the result $T^* \approx 0.5T_0$. First, consistent with the fact that the transition is locally driven, the transition temperature is proportional to the Kondo scale T_0 , the temperature characteristic of the local dynamics. Second, $T \approx 0.5T_0$ is also the temperature at which the slope of the real part of the self energy becomes one [12], so that the quasiparticle renormalization factor diverges. This result suggests that when the system cannot form a Fermi liquid due to a large residual scattering rate (which also yields a large residual entropy), it forms a superconducting state to quench the entropy. The antiferromagnetic transition temperature, on the other hand, depends upon both J (through the intersite exchange) and upon T_0 (through moment screening and superexchange). As noted above, we cannot determine whether these states coexist without a detailed exploration within the ordered phases. At half filling, where the antiferromagnetism produces an insulating phase, clearly the superconductivity will be suppressed for $J \leq 0.75$.

Our results offer a number of routes to be explored for explaining the complex superconducting phase diagrams of UPt_3 and $\text{U}_{1-x}\text{Th}_x\text{Be}_{13}$: 1) **Competition between phases with a multi-point irreducible star.** As mentioned above, for $\Upsilon_s < 0$ in finite d , $\vec{q} \neq 0$ pairs are favored. In this simple case of a bipartite lattice, the zone corner has a one-point irreducible star. For lattices with frustration, such as hexagonal UPt_3 and face-centered cubic UBe_{13} , it is possible to produce multi-point irreducible stars for staggered order parameters which can then have multiple phases [19]. 2) **Competition Between Different $\vec{q}$ values.** As noted above, T^* , the lower bound for the first order superconducting transition temperature is independent of $\vec{q}$ in our calculations. Thus multiple phases may thus correspond to superconducting transitions with different $\vec{q}$ values. 3) **Possible Instability of Spin Triplet/Channel Triplet Pairing.** At yet lower temperatures than those identified in Fig. 4, we observe a sign change in the pair field susceptibility associated with spin triplet-channel triplet odd-frequency pairing. Hence it is possible that the competition between this triplet-triplet and the singlet-singlet odd frequency pairing may explain the complex phase diagrams.

In closing, we note that this superconducting transition can agree with experiment only if it is weakly first order; this is plausible given the rapid change in free energy curvature at the order parameter origin. Detailed investigations in the ordered phase will resolve this issue and whether any of the above scenarios can describe the complex phase diagrams of UPt_3 and $\text{U}_{1-x}\text{Th}_x\text{Be}_{13}$.

We would like to acknowledge useful discussions with F. Anders, A.V. Balatsky, W. Chung, A. Georges, B. Goodman, D. Hess, H.R. Krishna-murthy, M. Ma, A.J.

Millis, and W. Putikka. Jarrell and Pang would like to acknowledge the support of NSF grants DMR-9406678 and DMR-9357199. Cox acknowledges the support of the NSF under Grant No. DMR-9420920. Computer support was provided by the Ohio Supercomputer Center.

-
- [1] Some good recent reviews are: N. Grewe and F. Steglich, *Handbook on the Physics and Chemistry of Rare Earths*, Eds. K.A. Gschneidner, Jr. and L.L. Eyring (Elsevier, Amsterdam, 1991) Vol. 14, P. 343; D. W. Hess, P. S. Riseborough, J. L. Smith, *Encyclopedia of Applied Physics* Eds. G. L. Trigg (VCH Publishers Inc., NY), Vol. 7 (1993) p. 435; H.R. Ott, *J. Low Temp. Phys.* **95**, 95, (1994).
- [2] J.A. Sauls, *Adv. in Phys.* **43**, 113–141 (1994).
- [3] V.L. Berezinski, *JETP Lett.* **20**, 287 (1974); A.V. Balatsky and E. Abrahams, *Phys. Rev. B* **45**, 13125 (1992); E. Abrahams, *et al.*, *Phys. Rev.* **B52**, 1271 (1995).
- [4] P. Coleman, E. Miranda, and A. Tsvetik, *Phys. Rev. Lett.*, **70**, 2960, (1993); P. Coleman, E. Miranda, and A.M. Tsvetik, *Phys. Rev. B* **49**, 8955 (1994).
- [5] V.J. Emery and S. Kivelson, *Phys. Rev. B* **46**, 10812 (1992).
- [6] A.W.W. Ludwig and I. Affleck, 1994, *Nucl. Phys. B* **428**, 545.
- [7] Enhanced staggered odd-frequency pairing correlations exist in the one-dimensional Kondo lattice when left and right movers approximately decouple, rendering effectively a two-channel lattice; see O. Zachar, S. Kivelson, and V. Emery, *Phys. Rev. Lett.* **77** 1342 (1996).
- [8] D.L. Cox, *Phys. Rev. Lett.* **59**, 1240 (1987); D.L. Cox, *Physica C* 153–155 1642 (1988); D.L. Cox, *Physica B* **186-188** (1993) 312. D.L. Cox and M.B. Maple, *Physics Today* **48**, 32 (1995); D.L. Cox and M. Jarrell, unpublished.
- [9] F. Steglich *et al.*, *Physica B* **223&224** (1996) 1.
- [10] This was noted much earlier in H.R. Ott, *Progress in Low Temperature Physics* **6**, 21 5 (1987).
- [11] P. Nozières, P. and A. Blandin, *J. de Phys. (Paris)* **41** (1980) 193.
- [12] M. Jarrell, H.-B. Pang, D.L. Cox, and K.-H. Luk, *Phys. Rev. Lett.* **77**, 1612 (1996); M. Jarrell, H.-B. Pang, and D.L. Cox, unpublished ; F. Anders, M. Jarrell, and D.L. Cox unpublished.
- [13] W. Metzner and D. Vollhardt, *Phys. Rev. Lett.* **62**, 324 (1989); see also E. Müller-Hartmann, *Z. Phys. B* **74**, 507 (1989).
- [14] Th. Pruschke, M. Jarrell and J.K. Freericks, *Adv. in Phys.* **42**, 187 (1995), and A. Georges, G. Kotliar, W. Krauth, and M. Rozenberg, *Rev. Mod. Phys.* **68**, 13 (Jan. 1996).
- [15] R.M. Fye and J.E. Hirsch, *Phys. Rev. B* **38**, 433 (1988).
- [16] K.-H. Luk, Mark Jarrell, and D.L. Cox, *Phys. Rev. B*, **50**, 15864 (1994).
- [17] M. Jarrell, *Phys. Rev. B*, **51**, 7429 (1995).
- [18] C. Owen and D. J. Scalapino, *Physica* **55**, 691(1971).

- [19] R. Heid, Ya.B. Bazaliy, V. Martisovits, and D.L. Cox, *Phys. Rev. Lett.* **74** (1995) 2571.




FIG. 1. Particle-particle interchannel opposite spin diagrams which contribute to the pair-field susceptibility. Here 1 and 2 label the channel and $\uparrow$ and $\downarrow$ the spin.




FIG. 2. Full antiferromagnetic χ_{AF} and ferromagnetic χ_F susceptibilities of the two-channel Kondo lattice with $J = 0.75$ and $N = 1.0$. χ_{AF} shows very little enhancement until $T \approx T_N$. As shown in the inset, the Kondo spin contribution to the susceptibility $\chi_{AF}^f(T)$ displays a protracted scaling region. The solid line is a fit of $1/\chi_{AF}^f(T)$ to the form $a(T - T_N)^\gamma$. The large difference between χ_{AF} and $\chi_{AF}^f(T)$ as well as the reduced exponent $\gamma < 1$ indicate the importance of screening of the local spin in this transition.




FIG. 3. Odd-frequency (spin-singlet, channel-singlet) pair-field susceptibility at both the zone center and corner. At low temperatures $T = T^* \leq T_c$ $P_{SsCs}(T)$ becomes negative, indicating a thermodynamic instability to the formation of a pairing state. The lack of a divergence of $P_{SsCs}(T)$ rules out a continuous transition, hence the system must undergo a discontinuous transition. In the inset, the dominant eigenvector of the pairing matrix M_- is plotted versus Matsubara frequency. It can be fit to the form $T/2\omega_n$ (solid line), which corresponds to the frequency variation of the gap function found in Ref.4




FIG. 4. Phase diagrams of the two-channel Kondo lattice for various values of J . The solid lines are fits to the data. For each value of J shown the antiferromagnetic transition occurs first (at a higher temperature) at half filling $N = 1$ (albeit very weakly when $J = 0.75$). T_N falls quickly with doping $N < 1$, with the transition becoming incommensurate near $T_N \rightarrow 0$. Away from half filling, superconductivity occurs first above a temperature $T^* \geq T_c$ is roughly equal to CT_0 , ($C = 0.43, 0.51$, and 0.58 , for $J = 0.5, 0.625, 0.75$ respectively) consistent with the local nature of the transition. T^* should be regarded as a lower bound on the superconducting transition temperature T_c (see text).